

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

PrTIMOPTIC-XE®

timolol maleate

sterile ophthalmic gellan solution, timolol, 0.5% w/v, ophthalmic

Elevated Intraocular Pressure Therapy

Elvium Life Sciences
575 Granite Court
Pickering, ON
L1W 3W8

Date of Initial Approval:
Oct. 28, 2016

Date of Revision:
August 19, 2020

Submission Control No: 241453

TIMOPTIC-XE® is a trademark of Elvium Life Sciences

RECENT MAJOR LABEL CHANGES

Not applicable

TABLE OF CONTENTS

TABLE OF CONTENTS	2
PART I: HEALTH PROFESSIONAL INFORMATION	3
1 INDICATIONS	3
1.1 Pediatrics.....	3
2 CONTRAINDICATIONS	3
3 DOSAGE AND ADMINISTRATION	3
3.1 Recommended Dose and Dosage Adjustment	3
3.2 Missed Dose.....	4
4 OVERDOSAGE	4
5 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	4
6 WARNINGS AND PRECAUTIONS	5
6.1 Special Populations	7
6.1.1 Pregnant Women	7
6.1.2 Breast-feeding	7
6.1.3 Pediatrics	7
7 ADVERSE REACTIONS	8
7.1 Clinical Trial Adverse Reactions.....	8
8 DRUG INTERACTIONS	9
8.1 Drug-Drug Interactions	9
8.2 Drug-Herb Interactions	10
8.3 Drug-Laboratory Test Interactions.....	10
9 ACTION AND CLINICAL PHARMACOLOGY	10
9.1 Mechanism of Action	10
9.2 Pharmacodynamics	10
9.3 Pharmacokinetics	11
10 STORAGE, STABILITY AND DISPOSAL	11
11 PHARMACEUTICAL INFORMATION	12
12 CLINICAL TRIALS	12
12.1 Study Results.....	12
13 NON-CLINICAL TOXICOLOGY	13
PATIENT MEDICATION INFORMATION	Error! Bookmark not defined.

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

TIMOPTIC-XE® (timolol maleate ophthalmic gellan solution) is indicated for:

- the reduction of elevated intraocular pressure.

1.1 Pediatrics

Pediatrics (<18 years of age): Safety and effectiveness in children have not been established.

2 CONTRAINDICATIONS

Timolol maleate is contraindicated in patients who are hypersensitive to this drug or to any ingredient in the formulation, including any non-medicinal ingredient, or component of the container. For a complete listing, see DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING.

- Reactive airway disease including bronchial asthma or a history of bronchial asthma, severe chronic obstructive pulmonary disease.
- Sinus bradycardia; sick sinus syndrome; sino-atrial block; second-and third-degree atrioventricular block; overt cardiac failure; cardiogenic shock.

3 DOSAGE AND ADMINISTRATION

3.1 Recommended Dose and Dosage Adjustment

Adults (≥18 years of age): One drop of TIMOPTIC-XE in the affected eye(s) once a day. Invert the closed container and shake once before each use. It is not necessary to shake the container more than once.

If needed, concomitant therapy with miotics, epinephrine and systemically administered carbonic anhydrase inhibitors may be given with TIMOPTIC-XE. Other topically applied medications should be administered no less than 10 minutes before TIMOPTIC-XE.

When using nasolacrimal occlusion or closing the eyelids for 2 minutes, the systemic absorption is reduced. This may result in an increase in local activity.

How to Transfer Patients from Other Therapy

When a patient is transferred from TIMOPTIC (timolol maleate ophthalmic solution, USP) to TIMOPTIC-XE, TIMOPTIC should be discontinued after proper dosing on one day, and treatment with the same concentration of TIMOPTIC-XE started on the following day.

When a patient is transferred from another topical ophthalmic beta-adrenergic blocking agent, that agent should be discontinued after proper dosing on one day and treatment with TIMOPTIC-XE started on the following day with one drop of TIMOPTIC-XE once a day.

When a patient is transferred from a single antiglaucoma agent, other than a topical ophthalmic beta-adrenergic blocking agent, continue the agent and add one drop of TIMOPTIC-XE to each affected eye once a day. On the following day, discontinue the previously used antiglaucoma agent and continue TIMOPTIC-XE.

Pediatrics (<18 years of age): Health Canada has not authorized an indication for pediatric use.

3.2 Missed Dose

If a dose is missed, it should be applied as soon as possible. However, if it is almost time for the next dose, the missed dose should be skipped and the next dose should be taken as usual.

4 OVERDOSAGE

For management of a suspected drug overdose, contact your regional poison control centre.

There have been reports of inadvertent overdose with TIMOPTIC (timolol maleate ophthalmic solution, USP) resulting in systemic effects similar to those seen with systemic beta-adrenergic blocking agents such as dizziness, headache, shortness of breath, bradycardia, bronchospasm, and cardiac arrest (see also ADVERSE REACTIONS).

The following additional therapeutic measures should be considered:

Gastric lavage: If ingested. Studies have shown that timolol does not dialyze readily.

Symptomatic bradycardia: Use atropine sulfate intravenously in a dosage of 0.25 to 2 mg to induce vagal blockade. If bradycardia persists, intravenous isoproterenol hydrochloride should be administered cautiously. In refractory cases the use of a transvenous cardiac pacemaker may be considered.

Hypotension: Use sympathomimetic pressor drug therapy, such as dopamine, dobutamine or levarterenol. In refractory cases the use of glucagon hydrochloride has been reported to be useful.

Bronchospasm: Use isoproterenol hydrochloride. Additional therapy with aminophylline may be considered.

Acute cardiac failure: Conventional therapy with digitalis, diuretics and oxygen should be instituted immediately. In refractory cases the use of intravenous aminophylline is suggested. This may be followed if necessary by glucagon hydrochloride which has been reported to be useful.

Heart block (second-or third-degree): Use isoproterenol hydrochloride or a transvenous cardiac pacemaker.

5 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table 1 – Dosage Forms, Strengths, Composition and Packaging.

Route of Administration	Dosage Form / Strength/Composition	Non-medicinal Ingredients
Ophthalmic	Solution / 0.5% w/v	Benzododecinium bromide 0.012% as preservative, gellan gum, mannitol, tromethamine and water for injection.

TIMOPTIC-XE is a formulation of timolol maleate containing novel delivery vehicle. It is a sterile aqueous solution. Each mL of TIMOPTIC-XE 0.5% w/v contains 5.0 mg of timolol (6.8 mg of timolol maleate).

Gellan gum used in this formulation contains a highly purified anionic heteropolysaccharide. Aqueous solutions of gellan gum form a clear transparent gel at low polymer concentrations in the presence of cations. The concentration of sodium cation in tears is ideally suited to cause gelation of the material when topically instilled in the conjunctival sac. When TIMOPTIC-XE contacts the pre-corneal tear film, it becomes a gel. The vehicle of TIMOPTIC-XE, gellan gum, increases the contact time of the drug with the eye.

TIMOPTIC-XE is a sterile, colourless to nearly colourless, slightly opalescent, slightly viscous, aqueous ophthalmic solution, supplied in translucent, high-density polyethylene ophthalmic dispenser, with a sealed dropper tip, a flexible fluted side area which is depressed to dispense the drops and a 2-piece cap assembly.

The opaque, white, 2-piece cap mechanism punctures the dropper tip seal upon initial use, then locks to provide a single cap during the usage period. Tamper evidence is provided by a safety strip on the container label.

Ophthalmic Gellan Solution TIMOPTIC-XE, equivalent to 5.0 mg (0.5% w/v) timolol per mL; in 5 mL dispensers, colour-coded with a dark blue label on the cap.

6 WARNINGS AND PRECAUTIONS

General

As with other topically applied ophthalmic drugs, this drug may be absorbed systemically. The same types of cardiovascular, pulmonary and other adverse reactions reported with systemic beta-adrenergic blocking agents may occur with topical administration.

Cardiovascular

In patients with cardiovascular diseases (e.g. coronary heart disease, Prinzmetal's angina and cardiac failure) and hypotension, therapy with beta-blockers should be critically assessed and the therapy with other active substances should be considered.

Cardiac failure should be adequately controlled before beginning therapy with TIMOPTIC-XE. Patients with a history of cardiovascular disease, including cardiac failure, should be watched for signs of deterioration of these diseases, and pulse rates should be monitored.

Due to its negative effect on conduction time, beta blockers should only be given with caution to patients with first degree heart block.

Respiratory reactions and cardiac reactions, including death due to bronchospasm in patients

with asthma and rarely death in association with cardiac failure, have been reported following administration of TIMOPTIC (timolol maleate ophthalmic solution, USP). These are also potential complications of therapy with TIMOPTIC-XE.

Because of the potential effects of beta-adrenergic blocking agents on blood pressure and pulse, these agents should be used with caution in patient with cerebrovascular insufficiency. If signs or symptoms suggesting reduced cerebral blood flow develop following initiation of therapy with TIMOPTIC-XE, alternative therapy should be considered.

Patients with severe peripheral circulatory disturbance/disorders (e.g. severe forms of Raynaud's disease or Raynaud's syndrome) should be treated with caution.

Endocrine and Metabolism

Masking of Hypoglycemic Symptoms in Patients with Diabetes Mellitus

TIMOPTIC-XE should be used with caution in patients subject to spontaneous hypoglycemia or in diabetic patients (especially those with labile diabetes) who are receiving insulin or oral hypoglycemic agents. Beta-adrenergic blocking agents may mask the signs and symptoms of acute hypoglycemia.

Masking of Thyrotoxicosis

β -adrenergic blocking agents may mask certain clinical signs of hyperthyroidism (e.g., tachycardia). Patients suspected of developing thyrotoxicosis should be managed carefully to avoid abrupt withdrawal of β -adrenergic blocking agents which might precipitate a thyroid storm.

Immune

Risk from Anaphylactic Reaction

While taking beta blockers, patients with a history of atopy or a history of severe anaphylactic reaction to a variety of allergens may be more reactive to repeated challenge with such allergens, either accidental, diagnostic, or therapeutic. These patients may be unresponsive to treatment of anaphylactic reactions with the usual doses of epinephrine since timolol may blunt the beta agonist effect of epinephrine. In such cases, alternatives to epinephrine should be considered.

Neurologic

Muscle Weakness

β -adrenergic blockade has been reported to increase muscle weakness consistent with certain myasthenic symptoms (e.g., diplopia, ptosis, and generalized weakness). Timolol has been reported rarely to increase muscle weakness in some patients with myasthenic symptoms.

Ophthalmologic

In patients with angle-closure glaucoma, the immediate objective of treatment is to reopen the angle. This requires constricting the pupil with a miotic. Timolol maleate has little or no effect on the pupil. When TIMOPTIC-XE is used to reduce elevated intraocular pressure in angle-closure glaucoma they should be used with a miotic and not alone.

Choroidal Detachment

Choroidal detachment has been reported with administration of aqueous suppressant therapy (e.g., timolol, acetazolamide or combination) after filtration procedures. Management of eyes with chronic or recurrent choroidal detachment should include stopping all forms of aqueous suppressant therapy and treating endogenous inflammation vigorously.

As with the use of other antiglaucoma drugs, diminished responsiveness to TIMOPTIC (timolol

maleate ophthalmic solution, USP) after prolonged therapy has been reported in some patients. However, in clinical studies in which 164 patients have been followed for at least 3 years, no significant difference in mean intra ocular pressure has been observed after initial stabilization.

Contact Lenses

The preservative in TIMOPTIC-XE is benzododecinium bromide. This preservative is a quaternary ammonium compound that may be absorbed by soft contact lenses. For TIMOPTIC-XE, studies have not been done in patients wearing contact lenses. However, in a clinical study, the time required to eliminate 50% of the gellan solution from the eye was up to 30 minutes.

Peri-Operative Considerations

Major Surgery

The necessity or desirability of withdrawal of β -adrenergic blocking agents prior to major surgery is controversial. Beta-adrenergic receptor blockade impairs the ability of the heart to respond to beta-adrenergically mediated reflex stimuli. This may augment the risk of general anesthesia in surgical procedures.

β -blocking ophthalmological preparations may block systemic β -agonist effects e.g. of adrenaline. The anaesthesiologist should be informed when the patient is receiving TIMOPTIC-XE.

Some patients receiving beta-adrenergic blocking agents have experienced protracted severe hypotension during anesthesia. Difficulty in restarting and maintaining the heartbeat has also been reported. For these reasons, in patients undergoing elective surgery, some authorities recommend gradual withdrawal of beta-adrenergic blocking agents. If necessary during surgery, the effects of β -adrenergic blocking agents may be reversed by sufficient doses of such agonists as isoproterenol, dopamine, dobutamine or levarterenol.

Respiratory

In patients with mild/moderate chronic obstructive pulmonary disease (COPD), TIMOPTIC-XE should be used with caution, and only if the potential benefit outweighs the potential risk.

6.1 Special Populations

6.1.1 Pregnant Women

TIMOPTIC-XE has not been studied in human pregnancy. The use of TIMOPTIC-XE requires that the anticipated benefit be weighed against possible hazards. Teratogenic studies in the mouse and rabbit at dose levels of 2 to 50 mg/kg/day did not reveal evidence of teratogenicity but did suggest embryotoxicity at the highest dose.

6.1.2 Breast-feeding

Timolol is detectable in human milk. Because of the potential for serious adverse reactions from timolol in nursing infants, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

6.1.3 Pediatrics

Pediatrics (<18 years of age): Safety and effectiveness in children have not been established.

7 ADVERSE REACTIONS

7.1 Clinical Trial Adverse Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

The most frequent (6.0%) drug-related complaint in the original clinical trials for TIMOPTIC-XE was transient blurred vision, lasting from 30 seconds to 5 minutes, following instillation.

Like other topically applied ophthalmic drugs, TIMOPTIC-XE is absorbed into the systemic circulation. This may cause similar undesirable effects as seen with systemic betablocking agents. Listed adverse reactions include reactions seen within the class of ophthalmic beta-blockers.

The following possibly, probably, or definitely drug-related adverse reactions occurred with frequency of at least 1% in active treatment controlled clinical trials:

Ocular

Burning and stinging, discharge, foreign body sensation, itching.

The adverse reactions reported for TIMOPTIC (timolol maleate ophthalmic solution, USP) are potential adverse reactions for TIMOPTIC-XE. Please refer to the list below.

Body as a Whole

Headache, asthenia, fatigue.

Cardiovascular

Aggravation or precipitation of certain cardiovascular pulmonary and other disorders presumably related to effects of systemic beta blockade has been reported (see CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS). These include bradycardia, chest pain, arrhythmia, hypotension, syncope, heart block, cerebrovascular accident, cerebral ischemia, palpitation, atrioventricular block, cardiac arrest, cardiac failure, congestive heart failure, edema, claudication, Raynaud's phenomenon, cold hands and feet and in insulin-dependent diabetics masked symptoms of hypoglycemia have been reported rarely.

Digestive

Dysgeusia, nausea, diarrhea, dyspepsia, dry mouth, abdominal pain, vomiting.

Hypersensitivity

Signs and symptoms of allergic reactions including anaphylaxis, angioedema, pruritus, urticaria, localized and generalized rash.

Immunologic

Systemic lupus erythematosus.

Integumentary

Alopecia, psoriasiform rash or exacerbation of psoriasis.

Metabolism and nutrition disorders

Hypoglycemia.

Musculoskeletal

Myalgia.

Nervous System/Psychiatric

Depression, insomnia, nightmares, memory loss, increases in signs and symptoms of myasthenia gravis, dizziness, paresthesia.

Respiratory

Bronchospasm (predominantly in patients with pre-existing bronchospastic disease), respiratory failure, dyspnea, cough.

Special Senses

Signs and symptoms of ocular irritation: including burning and stinging, itching, tearing, redness conjunctivitis, conjunctival injection, blepharitis, keratitis, blurred vision, decreased corneal sensitivity, and dry eyes. Visual disturbances: including refractive changes (due to withdrawal of miotic therapy in some cases), diplopia, corneal erosion ptosis, and choroidal detachment following filtration surgery (see WARNINGS AND PRECAUTIONS), tinnitus.

Urogenital

Decreased libido, Peyronie's disease, sexual dysfunction.

Causal Relationship Unknown

The following adverse reactions have been reported but a causal relationship to therapy with TIMOPTIC has not been established: aphakic cystoid macular edema, nasal congestion, anorexia, CNS effects (e.g., behavioral changes including confusion, hallucinations, anxiety, disorientation, nervousness, somnolence, and other psychic disturbances), hypertension, retroperitoneal fibrosis and pseudopemphigoid.

Adverse reactions reported in clinical experience with systemic timolol maleate may be considered potential side effects of ophthalmic timolol maleate.

8 DRUG INTERACTIONS

8.1 Drug-Drug Interactions

Beta-Adrenergic Blockers: Patients who are already receiving a beta blocker systemically and who are given TIMOPTIC-XE should be observed for a potential additive effect on the intraocular pressure or on the known systemic effects of beta blockers (hypotension and/or bradycardia). The concomitant use of two topical beta-adrenergic blocking agents is not recommended.

Calcium Channel Blockers or Catecholamine-Depleting Drugs: The potential exists for additive effects and production of hypotension and/or marked bradycardia when TIMOPTIC-XE is administered together with an oral calcium channel blocker or beta-adrenergic blocking agents, catecholamine-depleting drugs such as reserpine, antiarrhythmics, digitalis glycosides, parasymphomimetics or guanethidine.

Clonidine: Oral β -adrenergic blocking agents may exacerbate the rebound hypertension which

can follow the withdrawal of clonidine. If the two drugs are coadministered, the β -adrenergic blocking agent should be withdrawn several days before the gradual withdrawal of clonidine. If replacing clonidine by β -blocker therapy, the introduction of β -adrenergic blocking agents should be delayed for several days after clonidine administration has stopped.

Epinephrine: Although timolol maleate used alone has little or no effect on pupil size, mydriasis resulting from concomitant therapy with epinephrine has been reported occasionally. The potential for mydriasis also exists from concomitant therapy with TIMOPTIC-XE and epinephrine.

Quinidine: Potentiated systemic beta blockade (e.g., decreased heart rate, depression) has been reported during combined treatment with CYP2D6 inhibitors (e.g., quinidine, SSRIs) and timolol.

No specific drug interaction studies have been performed with TIMOPTIC-XE.

8.2 Drug-Herb Interactions

Interactions with herbal products have not been established.

8.3 Drug-Laboratory Test Interactions

Clinically important changes in standard laboratory parameters were rarely associated with the administration of systemic timolol maleate. Slight increases in blood urea nitrogen, serum potassium, serum uric acid and triglycerides and slight decreases in hemoglobin, hematocrit, and HDL-cholesterol occurred, but were not progressive or associated with clinical manifestations.

9 ACTION AND CLINICAL PHARMACOLOGY

9.1 Mechanism of Action

Timolol maleate is a general beta-adrenergic receptor blocking agent that does not have significant intrinsic sympathomimetic, direct myocardial depressant, or local anesthetic (membrane-stabilizing) activity. Timolol maleate combines reversibly with a part of the cell membrane, the beta-adrenergic receptor, and thus inhibits the usual biologic response that would occur with stimulation of that receptor. This specific competitive antagonism blocks stimulation of the beta-adrenergic receptors by catecholamines having beta-adrenergic stimulating (agonist) activity, whether these originate from an endogenous or exogenous source. Reversal of this blockade can be accomplished by increasing the concentration of the agonist, which will restore the usual biologic response.

9.2 Pharmacodynamics

TIMOPTIC (timolol maleate ophthalmic solution, USP) reduces elevated and normal intraocular pressure whether or not associated with glaucoma. Elevated intraocular pressure is a major risk factor in the pathogenesis of glaucomatous visual field loss. The higher the level of intraocular pressure, the greater the likelihood of glaucomatous visual field loss and optic nerve damage.

Onset of action of timolol maleate is usually rapid, occurring approximately 20 minutes after topical application on the eye. Maximum reduction of intraocular pressure occurs in one to two

hours with TIMOPTIC whereas it occurs in two to four hours with TIMOPTIC-XE. Significant lowering of intraocular pressure has been maintained for as long as 24 hours with 0.25% or 0.5% Ophthalmic Solution TIMOPTIC twice a day and TIMOPTIC-XE once a day. Repeated observations over a period of three years indicate that the intraocular pressure-lowering effect of TIMOPTIC is well maintained.

Timolol maleate is a non-selective beta-adrenergic receptor blocking agent that does not have significant intrinsic sympathomimetic, direct myocardial depressant, or local anesthetic (membrane-stabilizing) activity.

The precise mechanism of action of timolol maleate in lowering intraocular pressure is not clearly established at this time, although a fluorescein study and tonography studies indicate that its predominant action may be related to reduced aqueous formation. However, in some studies a slight increase in outflow facility was also observed. Unlike miotics, timolol maleate reduces intraocular pressure with little or no effect on accommodation or pupil size. Thus, changes in visual acuity due to increased accommodation are uncommon, and dim or blurred vision and night blindness produced by miotics are not evident. In addition, in patients with cataracts the inability to see around lenticular opacities when the pupil is constricted by miotics is avoided. When changing patients from miotics to TIMOPTIC a refraction might be necessary when these effects of the miotic have passed.

9.3 Pharmacokinetics

Absorption: Timolol maleate (S(-) enantiomer) is significantly metabolized after oral and ophthalmic administration.

Distribution: Timolol is moderately (<60%) bound to plasma proteins.

Metabolism: Based on correlation with debrisoquine metabolism, timolol metabolism is mediated primarily by cytochrome P-450 2D6.

Elimination: The drug and the metabolites (hydroxyethylamino, hydroxyethylglycolamino derivatives and a third minor metabolite that results from the hydroxylation of a terminal methyl group on the tertiary butylamino moiety) are excreted primarily via the kidney.

In a study of plasma drug concentration in six subjects, the systemic exposure to timolol was determined following twice-daily topical administration of timolol maleate ophthalmic solution 0.5% for 8 days. The mean peak plasma concentration following morning dosing was 0.46 ng/mL and following afternoon dosing was 0.35 ng/mL.

By comparison to plasma concentrations (10 to 20 ng/mL) following oral 5 mg dose, it was estimated that timolol was approximately 50% bio-available systemically following intraocular administration.

10 STORAGE, STABILITY AND DISPOSAL

Store at room temperature (15°-25°C). Protect from light and freezing.

The contents of TIMOPTIC-XE should not be used for more than one month after the date on which the container is first opened.

PRECAUTIONS). At trough (24 hours post-dose TIMOPTIC-XE, 12 hours post-dose TIMOPTIC), the mean reduction was 0.8 beats/minute for TIMOPTIC-XE and 3.6 beats/minute for TIMOPTIC; whereas at two hours post-dose, the mean reduction in heart rate was comparable (3.8 beats/minute for TIMOPTIC-XE and 5 beats/minute for TIMOPTIC).

There was a higher incidence of transient blurred vision following instillation in patients administered TIMOPTIC-XE.

In clinical trials TIMOPTIC-XE has been shown to reduce intraocular pressure in:

- Patients with chronic open-angle glaucoma
- Patients with ocular hypertension
- Aphakic patients having glaucoma, including those wearing contact lenses
- Patients with narrow angles and a history of spontaneous or iatrogenically-induced narrow-angle closure in the opposite eye in whom reduction of intraocular pressure is necessary (see WARNINGS AND PRECAUTIONS).

13 NON-CLINICAL TOXICOLOGY

Ocular Effects

No adverse ocular effects were observed in rabbits and dogs administered TIMOPTIC (timolol maleate ophthalmic solution, USP) topically in studies lasting one and two years respectively.

No adverse ocular effects were observed in rabbits and monkeys administered TIMOPTIC-XE topically in studies lasting one month and 12 months, respectively.

Table 2 - Acute Toxicity (LD₅₀)

Species and Age	Sex	Route of Administration	LD ₅₀ mg/kg
Mouse (A)	F	Oral	1190
	F	Intravenous	222
	F	Subcutaneous	1040
Rat (YA)	M	Oral	947
	F	Oral	900
	M	Oral (Fed)	1800
	M	Intraperitoneal	390
	F	Intraperitoneal	383
Rat (W)	M	Oral	1040
	F	Oral	969
	M/F	Intraperitoneal	409
Rat (I)	M/F	Oral	241
	M/F	Subcutaneous	143
Rabbit (A)	M/F	Oral	485
	M/F	Subcutaneous	34

(A)=Adult; (YA)=Young Adult; (W)=Weanling; (I)=Infant

Signs of toxicity occurred immediately after intravenous administration and from 10 to 30 minutes following oral, intraperitoneal or subcutaneous administration. The signs observed included lacrimation, ataxia, tremors and bradypnea. Clonic convulsions usually preceded death.

The oral LD50 of gellan gum is greater than 5 000 mg/kg in rats.

Oral Interactions Studies

Oral acute interaction studies in mice in which timolol maleate was administered with probenecid, methyldopa, hydralazine, hydrochlorothiazide, or tolbutamide, showed that these drugs had no influence on the toxicity of timolol maleate. Timolol maleate had no effect on the hypoprothrombinemia induced by bishydroxycoumarin in the dog.

Subacute Toxicity

In rats treated with 100 to 400 mg/kg/day for seven weeks, excessive salivation seen 5 to 10 minutes after dosing had a dose related incidence in the first week of the study. At necropsy, organ weight studies revealed a significant increase in the kidneys, spleen and liver of some treated animals. Except for splenic congestion, there were no morphological changes to account for the increase in organ weights. Rats treated with 1 gram per day for eight weeks exhibited ptyalism, muscle tremors and transient pale extremities.

In dogs, doses of 200 mg/kg/day or higher, were lethal to some animals. Low grade tubular nephrosis and trace amounts of hyaline casts in the collecting and convoluted tubules occurred in one of two dogs administered 100 mg/kg/day and in both dogs receiving 400 mg/kg/day. Small foci of tubular degeneration and regeneration occurred in the nephrotic areas. Similar slight multi focal degeneration of the collecting tubules in the medulla of both kidneys was evident in one of four dogs in a 15-day intravenous toxicity study.

Chronic Toxicity

Rats

Timolol was administered orally to rats at dose levels of 5, 10 and 25 mg/kg/day for up to 67 weeks. No physical signs, ocular signs or deaths which could be attributed to the drug were evident.

Dogs

In a 54-week oral study timolol was administered at doses of 5, 10 and 25 mg/kg/day. Body weight and food consumption were normal and no physical signs attributable to treatment were evident. Slight focal hyperplasia of the transitional epithelium was seen in the renal pelvis of one dog receiving 25 mg/kg/day.

Tumorigenic Tests

Lifetime studies with timolol have been completed in rats at oral doses of 25, 100 and 300 mg/kg/day and in mice at oral doses of 5, 50 and 500 mg/kg/day. In male and female rats and male mice at all dose levels, and in female mice at dose levels of 5 and 50 mg/kg/day, timolol demonstrated no carcinogenic effect. There was a slight increase in the incidence of mammary adenocarcinomas in female mice that received 500 mg/kg/day (about 500 times the maximum recommended human oral dose, on a mg/kg basis). Timolol caused dose-related elevations of serum prolactin in female mice at doses of 100 mg/kg/day or more, but only very slight transient elevations were found in male mice at doses of 500 mg/kg/day. Since numerous studies have demonstrated that drugs which cause elevations of serum prolactin are associated with mammary tumors in rodents, the mammary tumors in the female mice in the highest dosage group of this study were considered to have resulted from an increased serum prolactin. In

humans, no such association between serum prolactin and mammary carcinoma has been established.

Furthermore, in adult human female subjects who received oral dosages of up to 60 mg of timolol, the maximum recommended human oral dosage, there were no clinically meaningful changes in serum prolactin.

Reproductive Studies

Teratogenic studies in the mouse and rabbit at dose levels of 2 to 50 mg/kg/day did not reveal evidence of teratogenicity but did suggest embryotoxicity at the highest dose. Oral administration of timolol maleate to rats at dose levels of 4 to 100 mg/kg/day did not adversely affect the fertility of male or female rats, their reproductive performance, or the development of their offspring.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE
PATIENT MEDICATION INFORMATION

ⓇTIMOPTIC-XE®
Timolol maleate ophthalmic gellan solution

Read this carefully before you start taking **TIMOPTIC-XE** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **TIMOPTIC-XE**.

What is TIMOPTIC-XE used for?

TIMOPTIC-XE lowers the pressure in the eye for conditions such as open-angle glaucoma and ocular hypertension.

How does TIMOPTIC-XE work?

The active ingredient timolol maleate is a beta-blocking drug. It helps lower the pressure in the eye.

What are the ingredients in TIMOPTIC-XE?

Medicinal ingredients: timolol maleate.

Non-medicinal ingredients: gellan gum, mannitol, tromethamine and water for injection.
Benzododecinium bromide is added as preservative.

TIMOPTIC-XE comes in the following dosage forms:

TIMOPTIC-XE is a sterile eye drop containing timolol as the active ingredient. This eye drop also contains a new delivery vehicle which is obtained from a natural substance. This vehicle allows the eye drop to form a clear transparent gel when it contacts the eye.

Ophthalmic Gellan Solution TIMOPTIC-XE, equivalent to 5.0 mg (0.5%) timolol per mL; in 2.5 mL, 5 mL and 7.5 mL dispensers, colour-coded with a dark blue label on the cap.

Do not use TIMOPTIC-XE if:

- you are allergic to any of its components (see **What the ingredients in TIMOPTIC-XE** are section above);
- you have now or have had certain serious breathing problems such as asthma or chronic obstructive lung disease;
- you have certain heart diseases or conditions (such as slow or irregular heartbeats);

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take TIMOPTIC-XE. Talk about any health conditions or problems you may have, including:

- If you have any medical problems now or have had any in the past, especially asthma and other lung problems, heart problems or poor blood circulation.
- If you have any allergies
- If you have now or have had in the past, thyroid problems.
- If you have or have had in the past, diabetes or other blood sugar problems.
- If you are planning major surgery, including eye surgery, as using TIMOPTIC-XE may

change the effects of some medicines during anesthesia.

- If you had past eye problems such as choroidal detachment.
- If you had problems or develop problems with blood flow to the brain.
- If you are pregnant or intend to become pregnant.
- If you are breast feeding or intend to breast feed. Timolol has been detected in human breast milk.

TIMOPTIC-XE IS NOT RECOMMENDED FOR CHILDREN.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with TIMOPTIC-XE:

- Beta-blockers such as atenolol, epinephrine, quinidine,
- Calcium channel blockers or catecholamine depleting drugs such as reserpine.
- Clonidine
- Epinephrine

How to take TIMOPTIC-XE:

Read the following information carefully. **If you need any explanations, or further information, ask your physician or pharmacist.**

- Do not start taking any other medicines unless you have discussed the matter with your physician or pharmacist.
- If you develop any eye irritation or any new eye problems such as redness of the eye or swelling of the eyelids, contact your physician immediately.
- If you are using TIMOPTIC-XE with another eye drop, the drops should be instilled at least 10 minutes apart.
- Do not change the dosage recommended by your physician. If you must stop treatment, contact your physician immediately.
- Do not allow the tip of the container to touch the eye or areas around the eye. It may become contaminated with bacteria that can cause eye infections leading to serious damage of the eye, even loss of vision. To avoid possible contamination of the container, keep the tip of the container away from contact with any surface.
- The contents of the container should not be used for more than one month after the date on which the container is first opened.
- Do not administer while wearing (soft) contact lenses. Remove lenses before application and reinsert no earlier than 15 minutes after use. TIMOPTIC-XE contains benzalkonium chloride as a preservative. This preservative may be absorbed by soft contact lenses.

Instructions for use:

Step 1 - Before using the medication for the first time, be sure the Safety Strip on the front of the bottle is unbroken. A gap between the bottle and the cap is normal for an unopened bottle.




Figure A

Step 2 - Tear off the Safety Strip to break the seal.




Figure B

Step 3 - Invert the closed bottle and shake ONCE before each use. (It is not necessary to shake the bottle more than once.)

Step 4 - To open the bottle, unscrew the cap by turning as indicated by the arrows on the top of the cap. When opening the bottle for the first time, do not pull the cap directly up and away from the bottle. Pulling the cap directly up will prevent your container from operating properly.




Figure C

Step 5 - Tilt your head back and pull your lower eyelid down slightly to form a pocket between your eyelid and your eye.




Figure D

Step 6 - Invert the bottle and press lightly with the thumb or index finger over the “Finger Push Area” (as shown) until a single drop is dispensed into the eye as directed by your physician.




Figure E

Step 7 - After using TIMOPTIC-XE, press a finger into the corner of your eye, by the nose (as shown) for 2 minutes. This helps keep TIMOPTIC-XE in your eye.




Figure F

Step 8 - If drop dispensing is difficult after opening for the first time, replace the cap on the bottle and tighten (DO NOT OVERTIGHTEN) and then remove by turning the cap in the opposite direction as indicated by the arrows on the top of the cap.

Step 9 - Repeat steps 5 & 6 with the other eye if instructed to do so by your physician.

Step 10 - Replace the cap by turning until it is firmly touching the bottle. Do not overtighten or you may damage the bottle and cap.

Step 11 - The dispenser tip is designed to provide a single drop; therefore, do NOT enlarge the hole of the dispenser tip.

Step 12 - After you have used all doses, there will be some TIMOPTIC-XE left in the bottle. You should not be concerned since an extra amount of TIMOPTIC-XE has been added and you will get the full amount of TIMOPTIC-XE that your physician prescribed. Do not attempt to remove excess medicine from the bottle.

Usual adult dose:

One drop in the affected eye(s) once a day. Your doctor may change your dose based on your condition.

Overdose:

If you put too many drops in your eye or swallow any of the contents of the bottle, you should contact your physician immediately.

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

It is important to apply TIMOPTIC-XE as prescribed by your physician. If you miss a dose, apply it as soon as possible. However, if it is almost time for the next dose, skip the missed dose and go back to your regular dosing schedule. Do not double dose.

What are possible side effects from using TIMOPTIC-XE?

These are not all the possible side effects you may feel when taking TIMOPTIC-XE. If you experience any side effects not listed here, contact your healthcare professional.

You may experience:

- muscle pain
- abdominal pain
- nausea
- vomiting
- eye symptoms such as itching, burning and stinging, dry eyes, redness of the eye, foreign body sensation or visual changes, such as double vision.

Other side effects may also occur rarely, and some of these may be serious. These may include shortness of breath.

Timolol has been reported rarely to increase muscle weakness in some patients with myasthenia gravis.

Your vision may be temporarily blurred for 30 seconds to 5 minutes immediately after putting TIMOPTIC-XE in your eye. Make sure that your vision is clear before you drive a motor vehicle or operate machinery.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNCOMMON			
Slow heartbeat			✓
RARE			
Heart effects such as irregular heartbeat, heart block, low blood pressure			✓
Allergic reactions with symptoms such as swelling of the mouth and throat, shortness of breath, hives, severe itching and rash			✓

This is not a complete list of side effects. For any unexpected effects while taking TIMOPTIC-XE, contact your physician or pharmacist.

<p>Reporting Side Effects</p> <p>You can report any suspected side effects associated with the use of health products to Health Canada by:</p> <ul style="list-style-type: none"> • Visiting the Web page on Adverse Reaction Reporting (https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html) for information on how to report online, by mail or by fax; or • Calling toll-free at 1-866-234-2345. <p><i>NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.</i></p>
--

Storage:

- Store at room temperature 15°C - 25°C.
- Protect from light.
- Avoid freezing.
- Do not use this medicine for more than one month after the container was first opened.
- Keep out of the sight and reach of children.

If you want more information about TIMOPTIC-XE:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<http://hc-sc.gc.ca/index-eng.php>); the manufacturer's website <http://www.elvium.ca>, or by calling 1-833-744-0005.

This leaflet was prepared by Elvium Life Sciences.

Last Revised: August 19, 2020

TIMOPTIC-XE[®] is a trademark of Elvium Life Sciences.